

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-8020: Anti-Human CD194 (CCR4) (Mogamulizumab) - Biotin

Clone Name: KW-0761
Application: ELISA
Reactivity: Human

Alternative Name: CD194; CKR-4; CCR-4; CCR4; K5-5

Isotype: Human IgG1k

Immunogen Information: Humanization of mouse anti-CCR4 mAb7.

Description

Expression Host: HEK-293

This non-therapeutic biosimilar antibody uses the same variable region sequence as the therapeutic antibody Mogamulizumab. Clone KW-0761 recognizes human CD194 (CCR4). This product is for research use only.

Clone KW-0761 (Mogamulizumab) is a research-grade, afucosylated, humanized monoclonal antibody generated from mouse anti-CCR4 mAb7 that targets human CCR4.1 CC chemokine receptor type 4 (CCR4) is a protein that belongs to the G protein-coupled receptor family and is a receptor for a variety of CC chemokines including MCP-1, MIP-1, RANTES, TARC, and Macrophage-derived chemokine. Chemokines are involved in the development, homeostasis, and function of the immune system and are known to regulate cell trafficking of various types of leukocytes. In a 2018 Phase I clinical trial, Mogamulizumab was found to decrease the number of HTLV-1Â-infected cells and the levels of inflammatory markers related to HTLV-1Â-Associated Myelopathy.3

Product Info

Amount: $100 \mu g$

Concentration: 0.5 mg/ml

Content: This recombinant monoclonal antibody is aseptically packaged and formulated in 0.01 M

phosphate buffered saline (PBS) pH 7.2 - 7.4, 150 mM NaCl with no carrier protein, potassium,

calcium or preservatives added.

Functional grade biosimilar antibodies may be stored sterile as received at 2-8°C for up to one

Storage condition: month. For longer term storage, aseptically aliquot in working volumes without diluting and

store at -80°C. Avoid Repeated Freeze Thaw Cycles.

Application Note

The suggested concentration for Mogamulizumab biosimilar antibody for staining cells in flow cytometry is $\leq 1.0 \mu g$ per 106 cells in a volume of 100 μ l. Titration of the reagent is recommended for optimal performance for each application. ELISA