

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-8058: Anti-Human CTLA-4 (Ipilimumab) - Fc Muted™ Biotin

Clone Name: MDX-010
Application: ELISA
Reactivity: Human

Alternative Name: CD; GSE; GRD4; ALPS5; CD152; CTLA-4; IDDM12; CELIAC3

Isotype: Human IgG1k
Immunogen Information: Human CTLA-4

Description

Expression Host: HEK-293

This non-therapeutic biosimilar antibody uses the same variable region sequence as the therapeutic antibody Ipilimumab. Ipilimumab binds to Human CTLA-4. This product is for research use only.

Cytotoxic T-lymphocyteÂ-associated antigen 4 (CTLA-4) is a protein receptor that serves as an immune checkpoint and down-regulates the immune system. CTLA-4 is constitutively expressed in regulatory T cells but is only upregulated in conventional T cells following activation. Many cancers, including Melanoma, are associated with CTLA-4 upregulation because the bodyÂ's ability to recognize and destroy cancer cells is hampered by an inhibitory mechanism. Ipilimumab targets CTLA-4 and works by turning off this inhibitory mechanism and, thus, enhances the bodyÂ's own immune response against cancer cells.Â"2 Emerging research suggests that combined blockade of PD-1 and CTLA-4, with Nivolumab and Ipilimumab respectively, could produce greater antitumor activity than blockade of either pathway alone.1 This cost-effective, research-grade Anti-Human CTLA-4 (Ipilimumab) utilizes the same variable regions from the therapeutic antibody Ipilimumab making it ideal for research projects.

Product Info

Amount : 100 μg

Concentration: 0.5 mg/ml

Content: This Biotinylated antibody is formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS

pH 7.4, 1% BSA and 0.09% sodium azide as a preservative.

Storage condition : This biotinylated antibody is stable when stored at 2-8°C. Do not freeze.

Application Note

The suggested concentration for Ipilimumab biosimilar antibody for staining cells in flow cytometry is $<=1.0 \mu g$ per 106 cells in a volume of 100 μ l. Titration of the reagent is recommended for optimal performance for each application. ELISA