

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-8080: Anti-Human IL-2R alpha (CD25) (Basiliximab) - APC

Clonality: Monoclonal
Clone Name: Hu107
Application: FACS

**Alternative Name :** IL-2-RA; IL2-RA; TAC antigen; p55; CD25

**Isotype:** Human IgG1k **Immunogen Information:** Human CD25

## **Description**

Expression Host: HEK-293

This non-therapeutic biosimilar antibody uses the same variable region sequence as the therapeutic antibody Basiliximab. Basiliximab recognizes human CD25. This product is for research use only.

CD25, a 55 kD type I transmembrane glycoprotein, has been shown to play roles in lymphocyte differentiation, activation, and proliferation. Many resting memory T cells constitutively express IL2Ralpha. It functions as the receptor for HTLV-1, resulting in its expression on neoplastic cells in adult T cell lymphoma/leukemia. CD25 (sIL-2R) has been used to track disease progression. Some additional clinical applications include Chagas disease, a disease characterized by a decline of CD25 expression on immune cells, and Multiple sclerosis, in which treatments with mAbs target CD25. Anti-Human IL-2R alpha (CD25) (Basiliximab) utilizes the same variable regions from the therapeutic antibody Basiliximab making it ideal for research projects.

## **Product Info**

**Amount:** 50 μg

Concentration: 0.2 mg/ml

Content: This Allophycocyanin (APC) conjugate is formulated in 0.01 M phosphate buffered saline (150

mM NaCl) PBS pH 7.4, 1% BSA and 0.09% sodium azide as a preservative.

**Storage condition:** This Allophycocyanin (APC) conjugate is stable when stored at 2-8°C. Do not freeze.

## **Application Note**

The suggested concentration for Basiliximab biosimilar antibody for staining cells in flow cytometry is  $\leq 1.0 \, \mu g$  per 106 cells in a volume of 100  $\mu$ l. Titration of the reagent is recommended for optimal performance for each application.