9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

# 12-8087: Anti-Human IL-6 (Sarilumab) – HRP

| | |
|---|---|
| **Clonality :** | Monoclonal |
| **Clone Name :** | Hu137 |
| **Application :** | ELISA |
| **Reactivity :** | Human |
| **Alternative Name :** | Interleukin-6, CDF; HGF; HSF; BSF2; BSF-2; IFNB2; IFN-beta-2 |
| **Isotype :** | Human IgG1k |

## Description

Expression Host : HEK-293

This non-therapeutic antibody uses the same variable region sequence as the therapeutic antibody Sarilumab. Sarilumab binds to the transmembrane and soluble forms of the IL-6 receptor. This product is for research use only.
IL-6 and its signaling pathway play a part in immune response regulation, inflammation, and hematopoiesis.2 Sarilumab is a research-grade recombinant human monoclonal IL-6 receptor antagonist. It specifically binds to both the transmembrane and soluble forms of the IL-6 receptor, thus inhibiting IL-6Â–mediated cis and trans-signaling in a dose-dependent manner.1 Therapeutic Sarilumab, also known by the trade name Kevzara, is currently used to treat Rheumatoid Arthritis1, however, as of March 2020, The Feinstein Institute of Northwell Health publicized a study on "a human antibody that may prevent the activity" of IL-6 for the treatment of COVID-19.3 Anti-Human IL-6 (Sarilumab) utilizes the same variable regions from the therapeutic antibody Sarilumab making it ideal for research projects.

## Product Info

| | |
|---|---|
| **Amount :** | 100 µg |
| **Content :** | Concentration : 0.5 mg/ml<br>This HRP-conjugated antibody is formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4, 1% BSA. (Warning: Use of sodium azide as a preservative will inhibit the enzyme activity of horseradish peroxidase) |
| **Storage condition :** | This horseradish peroxidase conjugated monoclonal antibody is stable when stored at 2-8°C. Do not freeze. |

## Application Note

The suggested concentration for Alemtuzumab biosimilar antibody for staining cells in flow cytometry is <= 1.0 µg per 106 cells in a volume of 100 µl. Titration of the reagent is recommended for optimal performance for each application.ELISA

***For Research Use Only. Not for use in diagnostic/therapeutics procedures.***