

## 12-9270: Anti-DNAM1 antibody(DM96), Rabbit mAb

**Clonality :** Monoclonal  
**Clone Name :** DM96  
**Application :** ELISA,FACS  
**Reactivity :** Human  
**Alternative Name :** DNAM-1, CD226, PTA1

### Description

This gene encodes a glycoprotein expressed on the surface of NK cells, platelets, monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. The protein mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Alternative splicing results in multiple transcript variants.

### Product Info

**Amount :** 100 µg  
**Purification :** Purified from cell culture supernatant by affinity chromatography  
**Content :** Not Sterile  
**Storage condition :** Store at -20°C for 12 months (Avoid repeated freezing and thawing)

### Application Note

ELISA 1/5000-10000;FACS 1/100




Figure 1. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human DNAM-1 protein, mFc-His tagged protein can bind Rabbit anti-DNAM-1 monoclonal antibody (clone: DM96) in a linear range of 0.64-80 ng/ml.




Figure 2. Expi 293 cell line transfected with irrelevant protein (A) and human DNAM-1 (B) were surface stained with Rabbit anti-DNAM-1 monoclonal antibody 1 $\mu$ g/ml (clone: DM96) followed by PE-conjugated anti-rabbit IgG secondary antibody.