

## 12-9286: Anti-OX40-L antibody(DM112), Rabbit mAb

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	DM112
<b>Application :</b>	ELISA,FACS
<b>Reactivity :</b>	Human
<b>Alternative Name :</b>	OX40L, TNFSF4, CD252, Glycoprotein Gp34, TXGP1, CD134 ligand, CD134L

### Description

This gene encodes a cytokine of the tumor necrosis factor (TNF) ligand family. The encoded protein functions in T cell antigen-presenting cell (APC) interactions and mediates adhesion of activated T cells to endothelial cells. Polymorphisms in this gene have been associated with Sjogren's syndrome and systemic lupus erythematosus. Alternative splicing results in multiple transcript variants

### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Purified from cell culture supernatant by affinity chromatography
<b>Content :</b>	Not Sterile
<b>Storage condition :</b>	Store at -20°C for 12 months (Avoid repeated freezing and thawing)

### Application Note

ELISA 1/5000-10000;FACS 1/100




Figure 1. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human OX40L protein, mFc-His tagged protein can bind Rabbit anti-OX40L monoclonal antibody (clone: DM112) in a linear range of 0.2-90 ng/ml.




Figure 2. Flow cytometry analysis with Anti-OX40L (DM112) on Expi293 cells transfected with human OX40L (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).