

30-2745: Anti-Hu CLEC2 PE

| | |
|--------------------------------|---|
| Clonality : | Monoclonal |
| Clone Name : | AYP1 |
| Application : | FACS |
| Reactivity : | Human |
| Conjugate : | PE |
| Gene : | CLEC1B |
| Gene ID : | 51266 |
| Uniprot ID : | Q9P126 |
| Alternative Name : | C-type lectin domain family 1 member B CLEC1B, CLEC2B, PRO1384, QDED721 |
| Immunogen Information : | A recombinant extracellular domain of human CLEC2 (amino acids 68-229) |

Description

CLEC2 (C-type lectin-like receptor 2) functions as a platelet receptor for the lymphatic endothelial marker, PDPN, and mediates platelet activation. Besides platelets, it can be found on myeloid cells and NK cells. CLEC2 functions also as an attachment factor for HIV-1 and facilitates its capture by platelets. Platelet-aggregating snake venom protein rhodocytin also binds to CLEC2.

Specificity : The mouse monoclonal antibody AYP1 recognizes an epitope within the extracellular part of CLEC2, a transmembrane glycoprotein expressed on activated platelets and on platelet microparticles.

Product Info

| | |
|----------------------------|---|
| Amount : | 100 Tests |
| Purification : | Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography. |
| Content : | Storage Buffer: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |
| Storage condition : | Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze. |

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10⁵ cells / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.