

30-2759: Anti-Hu CD77 FITC

Clonality :	Monoclonal
Clone Name :	38.13
Application :	FACS
Reactivity :	Human
Conjugate :	FITC
Alternative Name :	globotriaosylceramide Gb3, Gal-alpha1-4Gal-beta1-4Glc-beta1-Cer, P blood group
Immunogen Information :	Daudi cell line (Burkitt's lymphoma)

Description

CD77 (globotriaosylceramide Gb3), also known as the Pk blood group antigen, BLA (Burkitt's lymphoma associated antigen), or CTH (ceramide trihexoside) is a neutral glycosphingolipid composed of three carbohydrate molecules linked to a lipid moiety in the cell membrane (Gal-alpha1-4Gal-beta1-4Glc-beta1-Cer). It is expressed on germinal center B cells, Burkitt's lymphoma cells, it can be induced on extrafollicular B cells and it is also found on endothelia and epithelia. CD77 may be involved in elimination of germinal center B cells that fail to produce high affinity antibodies, and serves also as receptor for shiga toxin and verotoxin.

Specificity : The rat monoclonal antibody 38.13 recognizes CD77 (globotriaosylceramide Gb3), a neutral glycosphingolipid expressed mainly on the surface of B cell populations, such as germinal center B cells and Burkitt's lymphoma cells.

Product Info

Amount :	100 Tests
Purification :	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Content :	Storage Buffer: Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Storage condition :	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.