

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

11-7017: Polyclonal Antibody to Histone H2A

Clonality: Polyclonal
Application: IHC,WB
Reactivity: Human
Gene: HIST1H2AB
Gene ID: 3012

Gene ID: 3012
Uniprot ID: P04908
Format: Purified

Alternative Name: HIST1H2AB,H2AFM

Isotype: Rabbit IgG

Immunogen Information: Full length recombinant Histone H2A protein was used as the immunogen for this antibody.

Description

Histone H2A is one of the five main histone proteins that is approximately 14 kDa protein consisting of 130 amino acid residues and present in the nucleosome in the form of H2A-H2B heterodimer. H2A is mainly composed of non-allelic variants and interestingly atypical variants, such as, MacroH2As, H2A.X, H2A-Bbd, differ considerably in size with H2A. H2A undergoes various post-translational modifications, including acetylation, phosphorylation, methylation, and ubiquitination, which occur in response to various stimuli and directly affect the accessibility of chromatin to transcription factors. This gene is mainly expressed on blood, brain, breast and it plays a significant role in the disease, riddle syndrome.

Product Info

Amount : 25 μg / 100 μg

Purification : Protein A Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western blot analysis: 2-4 μg/ml, Immunohistochemical analysis: 5 μg/ml




Fig-1: Western blot analysis of Histone H2A. Anti- Histone H2A antibody (11-7017) was used at 2 μ g/ml on Brain lysate.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-2: Immunohistochemical analysis of Histone H2A. Anti-Histone H2A antibody (11-7017) in human Liver tissue at 5 μ g/ml.




Fig-3: Immunohistochemical analysis of Histone H2A. Anti-Histone H2A antibody (11-7017) in human Lung tissue at 5 μ g/ml.