# **w** abeomics

### 36-1752: Monoclonal Antibody to TNF-alpha (Tumor Necrosis Factor alpha)(TNF706 + P/T2)

| Clonality :           | Monoclonal  |
|-----------------------|---|
| Clone Name :          | TNF706 + P/T2   |
| Application :         | IHC   |
| Reactivity :          | Human, Mouse, Rat   |
| Gene :                | TNF   |
| Gene ID :             | 7124  |
| Uniprot ID :          | P01375  |
| Format :              | Purified  |
| Alternative Name :    | TNF,TNFA,TNFSF2   |
| Isotype :             | Mouse IgM, kappa  |
| Immunogen Information | Recombinant N-terminal fragment of human TNF-? (TNF706); Peptide corresponding to aa115-130 of human TNF-? (P/T2) |

#### Description

TNF alpha is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF alpha causes cytolysis of certain transformed cells. Although it has little effect on many cultured normal human cells, TNF alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. TNF alpha is currently being evaluated in treatment of certain cancers and AIDS Related Complex.

#### **Product Info**


| Amount :            | 100 µg  |
|---------------------|---|
| Purification :      | Affinity Chromatography   |
| Content :           | 100 $\mu g$ in 500 $\mu l$ PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.                     |
| Storage condition : | Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles. |

#### **Application Note**

Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris Buffer with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),

## **₩** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Formalin-fixed, paraffin-embedded human Erdheim Chester disease (also known as polyostotic sclerosing histiocytosis) stained with TNF alpha Monoclonal Antibody (TNF706 + P/T2).