

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2811: Anti-Hu CD272 Purified

Clone Name: Monoclonal
Clone Name: MIH26
Application: FACS

Reactivity: Human, Non-human primates

 Gene :
 BTLA

 Gene ID :
 151888

 Uniprot ID :
 Q7Z6A9

 Format :
 Purified

Alternative Name: B and T lymphocyte associated BTLA1 Immunogen Information: cells transfected with human CD272

Description

CD272, a type I transmembrane glycoprotein, contains in its intracellular domain two ITIM sequences, which are upon CD272 triggering phosphorylated and recruit SHP phosphatases to attenuate cell activation. CD272 is expressed on B and T lymphocytes, NK cells, dendritic cells, and macrophages, and its ligand is CD270. Defects in CD272-CD270 inhibitory mechanism lead to autoimmune diseases. Overexpression of CD272 is a marker of tolerant T cells.

Specificity: The mouse monoclonal antibody MIH26 recognizes an extracellular epitope of CD272, a transmembrane glycoprotein serving as a negative regulator of the activation in various leukocyte types.

Product Info

Amount: 0.1 mg

Purification : Purified by protein-A affinity chromatography.

Content: Concentration: 1 mg/ml

Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow cytometry: Recommended dilution: 2-5 µg/ml.