

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-2829: Anti-Hu CD11b APC

Clone Name: Monoclonal ICRF44
Application: FACS

**Reactivity:** Human, Non-human primates

Conjugate: APC
Gene: ITGAM
Gene ID: 3684
Uniprot ID: P11215

Alternative Name: integrin subunit alpha M Mac-1, Integrin alpha M, ITGAM, CR3A, MO1A, MAC1A

Immunogen Information: Rheumatoid synovial cells and human monocytes.

## **Description**

CD11b (integrin alphaM subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with integrin beta2 subunit (CD18); expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/CD18 integrin (Mac-1, CR3) is highly expressed on NK cells, neutrophils, monocytes and less on macrophages. CD11b/CD18 integrin is implicated in various adhesive interactions of monocytes, macrophages and granulocytes, facilitating their diapedesis, as well as it mediates the uptake of complement coated particles, serving as a receptor for the iC3b fragment of the third complement component.

Specificity: The mouse monoclonal antibody ICRF44 recognizes an extracellular epitope of CD11b (Mac-1alpha), a 165-170 kDa type 1 transmembrane protein mainly expressed on monocytes, granulocytes and NK-cells.

## **Product Info**

Amount: 100 Tests

Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions

**Purification:** and unconjugated antibody and free fluorochrome are removed by size-exclusion

chromatography.

Content: Concentration: 0.5 mg/ml

Storage Buffer: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

**Storage condition :** Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

## **Application Note**

Flow cytometry: The reagent is designed for analysis of human blood cells using 10  $\tilde{A} \square \hat{A} \mu l$  reagent / 100  $\tilde{A} \square \hat{A} \mu l$  of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.