

## 30-2856: Anti-Human TCR gamma/delta (Clone: 11F2) FITC Conjugated

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	11F2
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	FITC
<b>Alternative Name :</b>	TCRG/D
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Purified CD3 antigen complex

### Description

Specificity: The mouse monoclonal antibody 11F2 recognizes an extracellular epitope in all molecular forms of the TCR gamma/delta.

The antigen-specific T cell receptor (TCR) is composed of either alpha and beta subunit, or gamma and delta subunit. Majority of T cells present in the blood, lymph and secondary lymphoid organs express TCR alpha/beta heterodimers, whereas the T cells expressing TCR gamma/delta heterodimers are localized mainly in epithelial tissues and at the sites of infection. The subunits of TCR heterodimers are covalently bonded and in the endoplasmic reticulum they associate with CD3 subunits to form functional TCR-CD3 complex. Lack of expression of any of the chains is sufficient to stop cell surface expression.

### Product Info

<b>Amount :</b>	100 tests
<b>Purification :</b>	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
<b>Content :</b>	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

### Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.




Fig1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human TCR gamma/delta (11F2) FITC antibody (4 1/4l reagent / 100 1/4l of peripheral whole blood).




Fig2: Flow cytometry multicolor surface staining of human lymphocytes stained using anti-human TCR gamma/delta (11F2) FITC antibody (4  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood) and anti-human CD3(UCHT1) APC antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).