

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-2913: Anti-Hu CLEC2 APC Mab (AYP1)

Clonality: Monoclonal

Clone Name : AYP1
Application : FACS
Reactivity : Human
Conjugate : APC
Gene : CLEC1B
Gene ID : 51266
Uniprot ID : Q9P126

Alternative Name: CLEC1B, CLEC2B, PRO1384, QDED721

**Isotype:** Mouse IgG1 kappa

Immunogen Information: A recombinant extracellular domain of human CLEC2 (amino acids 68-229)

## **Description**

Specificity: The mouse monoclonal antibody AYP1 recognizes an epitope within the extracellular part of CLEC2, a transmembrane glycoprotein expressed on activated plateles and on platelet microparticles.

CLEC2 (C-type lectin-like receptor 2) functions as a platelet receptor for the lymphatic endothelial marker, PDPN, and mediates platelet activation. Besides platelets, it can be found on myeloid cells and NK cells. CLEC2 functions also as an attachment factor for HIV-1 and facilitates its capture by platelets. Platelet-aggregating snake venom protein rhodocytin also binds to CLEC2.

## **Product Info**

Amount: 100 Tests

Purification: Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions

and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Content: Storage Buffer: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

**Storage condition :** Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

## **Application Note**

Flow cytometry: The reagent is designed for analysis of human blood cells using 10  $\mu$ l reagent / 100  $\mu$ l of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CLEC2 (AYP1) APC antibody (10  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).




Figure 2: