9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

# 32-17641: Recombinant Human CFB Protein, hFc Tag

**Uniprot ID :** P00751
**Alternative Name :** AHUS4, ARMD14, BF, BFD, CFAB, CFBD, FB, FBI12, GBG, H2-Bf, PBF2

## Description

Molecular Characterization: CFB?Thr26-Leu764?+hFc(Glu99-Ala330)
Molecular Weight: The protein has a predicted molecular mass of 109.2 kDa after removal of the signal peptide.The apparent molecular mass of CFB-hFc is approximately 100-130 kDa due to glycosylation.
Description: Recombinant Human CFB with C-terminal human Fc tag
This gene encodes complement factor B, a component of the alternative pathway of complement activation. Factor B circulates in the blood as a single chain polypeptide. Upon activation of the alternative pathway, it is cleaved by complement factor D yielding the noncatalytic chain Ba and the catalytic subunit Bb. The active subunit Bb is a serine protease which associates with C3b to form the alternative pathway C3 convertase. Bb is involved in the proliferation of preactivated B lymphocytes, while Ba inhibits their proliferation. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. This cluster includes several genes involved in regulation of the immune reaction. Polymorphisms in this gene are associated with a reduced risk of age-related macular degeneration. The polyadenylation site of this gene is 421 bp from the 5' end of the gene for complement component 2.

## Product Info

| | |
|---|---|
| **Amount :** | 100 µg / 50 µg |
| **Content :** | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. |
| **Storage condition :** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |

***For Research Use Only. Not for use in diagnostic/therapeutics procedures.***