

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-18231: Human VEGF121 Protein, hFc Tag

**Uniprot ID:** P15692-9

Alternative Name: MVCD1, VEGF, VPF, VEGFA

## **Description**

Description: Recombinant human VEGF121 Protein with C-terminal human Fc tag

Background: Vascular endothelial growth factor (VEGF) is also known as vascular permeability factor (VPF) and VEGF-A, and is a member of the platelet-derived growth factor (PDGF)/vascular endothelial growth factor (VEGF) family and encodes a protein that is often found as a disulfide linked homodimer. This protein is a glycosylated mitogen that specifically acts on endothelial cells and has various effects, including mediating increased vascular permeability, inducing angiogenesis, vasculogenesis and endothelial cell growth, promoting cell migration, and inhibiting apoptosis. Alternatively spliced transcript variants, encoding either freely secreted or cell-associated isoforms, have been characterized. Alternatively spliced isoforms of 121,145,165,183,189 and 206 amino acids in length are expressed in humans. VEGF165 appears to be the most abundant and potent isoform, followed by VEGF121 and VEGF189. VEGF121 is the only form that lacks a basic heparinbinding region and is freely diffusible. Mouse embryos expressing only the corresponding isoform (VEGF120) do not survive to term, and show defects in skeletogenesis. Human VEGF121 shares 87% as sequence identity with corresponding regions of mouse and rat, 93% with feline, equine and bovine, and 91%, 95% and 96% with ovine, canine and porcine VEGF, respectively. VEGF121 induces the proliferation of lymphatic endothelial cells. The lymphangiogenesis may be promoted by upregulation of VEGF121, which may in turn act in part via induction of VEGF-C.

Description: Recombinant human VEGF121 Protein with C-terminal human Fc tag

Molecular Characterization: VEGF121(Ala27-Arg147) hFc(Glu99-Ala330)

Molecular Weight :The protein has a predicted molecular mass of 40.2 kDa after removal of the signal peptide. The apparent molecular mass of VEGF121-hFc is approximately 35-55 kDa due to glycosylation.

Tag: C-Human Fc tag

## **Product Info**

Amount :  $50 \mu g / 100 \mu g$ 

Purification:

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

**Content:** Lyophilized from sterile PBS, pH 7.4. Normally 5 % -8% trehalose is added as protectants before

lyophilization.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

**Storage condition :** for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.