

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-18245: Human KIR2DL3 Protein, His Tag

Uniprot ID <sub>P43628</sub>

Name:

**Alternative** p58;NKAT;GL183;NKAT2;CD158b;KIR2DL;NKAT2A;NKAT2B;CD158B2;KIR-K7b;KIR-K7c;KIR2DS5;KIRCL23;KIR-023GB

## **Description**

Description: Recombinant Human KIR2DL3 Protein with C-terminal 6XHis tag

Background: Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response.

Molecular Characterization: mass of 25.4 kDa after removal of the signal peptide. The apparent molecular mass of KIR2DL3-His is approximately 35-55 kDa due to glycosylation.

Tag: C-6×His Tag

## **Product Info**

Amount: 50 μg / 100 μg

The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue Purification:

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before Content:

lyophilization.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for Storage condition:

use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized

proteins are shipped at ambient temperature.




Figure 1.Human KIR2DL3 Protein, His Tag on SDS-PAGE under reducing condition.