

32-18258: Human IL21R Protein, mFc Tag

Uniprot ID : Q9HBE5 Alternative Name : CD360;IMD56;NILR

Description

Description : Recombinant Human IL21R with C-terminal mouse Fc tag

Background : The protein encoded by this gene is a cytokine receptor for interleukin 21 (IL21). It belongs to the type I cytokine receptors, and has been shown to form a heterodimeric receptor complex with the common gamma-chain, a receptor subunit also shared by the receptors for interleukin 2, 4, 7, 9, and 15. This receptor transduces the growth promoting signal of IL21, and is important for the proliferation and differentiation of T cells, B cells, and natural killer (NK) cells. The ligand binding of this receptor leads to the activation of multiple downstream signaling molecules, including JAK1, JAK3, STAT1, and STAT3. Knockout studies of a similar gene in mouse suggest a role for this gene in regulating immunoglobulin production. Three alternatively spliced transcript variants have been described.

Molecular Characterization: mass of 50.7 kDa after removal of the signal peptide. The apparent molecular mass of IL21RmFc is approximately 55-70 kDa due to glycosylation.

Tag :C-Mouse Fc Tag

Product Info

| Amount : | 50 μg / 100 μg |
|---------------------|---|
| Purification : | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Content : | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. |
| Storage condition : | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |



Figure 1. Human IL21R Protein, mFc Tag on SDS-PAGE under reducing condition



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure 2. ELISA plate pre-coated by 2 \hat{l}_{4} g/mL (100 \hat{l}_{4} L/well) Human IL21R Protein, mFc Tag can bind Anti-IL21R antibody(DMC225), IgG1 Chimeric mAb in a linear range of 3.20-80 ng/mL.