

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-18454: Human HGF(1-728) Protein, hFc Tag

P14210 **Uniprot ID:**

Alternative Name: SF; HGFB; HPTA; F-TCF; DFNB39

Description

Description: Recombinant human HGF(1-728) Protein with C-terminal human Fc tag

Background: This gene encodes a protein that binds to the hepatocyte growth factor receptor to regulate cell growth, cell motility and morphogenesis in numerous cell and tissue types. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate alpha and beta chains, which form the mature heterodimer. This protein is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. This protein also plays a role in angiogenesis, tumorogenesis, and tissue regeneration. Although the encoded protein is a member of the peptidase S1 family of serine proteases, it lacks peptidase activity. Mutations in this gene are associated with nonsyndromic hearing loss.

Molecular Characterization: mass of 109.3 kDa after removal of the signal peptide. The apparent molecular mass of HGF(1-728)-hFc is approximately 70-130 kDa due to glycosylation.

Tag: C-Human Fc tag

Product Info

Amount: $50 \mu g / 100 \mu g$

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue Purification:

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before Content:

lyophilization.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for Storage condition:

use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized

proteins are shipped at ambient temperature.




Figure 1. Human HGF(1-728) Protein, hFc Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 μg/mL (100 μL/well) Human HGF(1-728) Protein, hFc Tag can bind Anti-HGF(ficlatuzumab biosimilar) mAb in a linear range of 16-2000 ng/mL.