

## 11-3025: Polyclonal Antibody to mTLR3

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	FACS, WB
<b>Reactivity :</b>	Mouse
<b>Gene :</b>	Tlr3
<b>Gene ID :</b>	142980
<b>Uniprot ID :</b>	Q99MB1
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Tlr3
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	A partial length recombinant mTLR3 protein (amino acids 180-400) was used as the immunogen for this antibody.

### Description

Mouse Toll-like receptor 3 (mTLR3) belongs to a family of evolutionary conserved innate immune recognition molecules and recognizes double-stranded RNA, a molecular pattern associated with viral infections. It has seven exons. mTlr3 show high homology to hTLR3 in their predicted cDNA and protein sequences. The intracellular region of mTlr3 has remarkable homology with the same region of others mTlrs. mTlr3 mRNA is ubiquitously expressed in tissues, being expressed highest in spleen, kidney and lung. Its activation by the synthetic ligand polyinosine:polycytidylic acid (poly I:C) or by mRNA rapidly causes growth cone collapse and irreversibly inhibits neurite extension independent of NF-kappaB. The high levels of mTlr3 expression in Mz B cells perhaps contribute to their ability to clear antigen from the blood.

### Product Info

<b>Amount :</b>	25 µg / 100 µg
<b>Purification :</b>	Protein A Chromatography
<b>Content :</b>	25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Western blot analysis: 2-4 µg/ml, Flowcytometric analysis: 1-2 µg/10<sup>6</sup> Cells




Fig-1: Western blot analysis of mTLR3. Anti- mTLR3 antibody (11-3025) was used at 2 µg/ml on EL-4 lysate.




Fig-2: Intracellular flowcytometric staining of mTLR3 in Splenocytes using 1 µg/10<sup>6</sup> cells of Anti-mTLR3 (11-3025). Green represent Isotype control and red represent Anti-mTLR3 antibody (Abeomics). Goat Anti-Rabbit IgG FITC conjugated was used as the secondary antibody.