

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-9426: Anti-Nectin4 antibody(DMC438); IgG1 Chimeric mAb

Clonality: Monoclonal
Clone Name: DMC438
Application: FACS
Reactivity: Human
Uniprot ID: Q96NY8

Alternative Name : EDSS1; LNIR; nectin-4; PRR4; PVRL4 **Isotype :** Rabbit/Human Fc chimeric IgG1

Description

Description :Anti-Nectin4 antibody(DMC438); IgG1 Chimeric mAb

This gene encodes a member of the nectin family. The encoded protein contains two immunoglobulin-like (Ig-like) C2-type domains and one Ig-like V-type domain. It is involved in cell adhesion through trans-homophilic and -heterophilic interactions. It is a single-pass type I membrane protein. The soluble form is produced by proteolytic cleavage at the cell surface by the metalloproteinase ADAM17:TACE. The secreted form is found in both breast tumor cell lines and breast tumor patients. Mutations in this gene are the cause of ectodermal dysplasia-syndactyly syndrome type 1; an autosomal recessive disorder. Alternatively spliced transcript variants have been found but the full-length nature of the variant has not been determined.

Product Info

Amount : $10 \mu g / 100 \mu g$

Purification: Purified from cell culture supernatant by affinity chromatography

Content: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

Storage condition : for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Application Note

FACS 1:100




Figure 1. Flow cytometry analysis with Anti-Nectin4 (DMC438) on Expi293 cells transfected with human Nectin4 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).