

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-9452: Anti-CD164 antibody(DMC476); IgG1 Chimeric mAb

Clonality: Monoclonal
Clone Name: DMC476
Application: FACS
Reactivity: Human
Uniprot ID: Q04900

Alternative Name: LMOR; M-OR-1; MOP; MOR; MOR1; OPRM

**Isotype:** Rabbit/Human Fc chimeric IgG1

## **Description**

Description: Anti-CD164 antibody(DMC476); IgG1 Chimeric mAb

This gene encodes one of at least three opioid receptors in humans; the mu opioid receptor (MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as beta-endorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse; such as nicotine; cocaine; and alcohol via its modulation of the dopamine system. The NM\_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6 transmembrane domains. [provided by RefSeq; Oct 2013]

## **Product Info**

Amount:  $10 \mu g / 100 \mu g$ 

**Purification:** Purified from cell culture supernatant by affinity chromatography

Content: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

**Storage condition:** for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

## **Application Note**

FACS 1:100




Figure 1. CD164 protein is highly expressed on the surface of Expi293 cell membrane. Flow cytometry analysis with Anti-CD164 (DMC476) on Expi293 cells transfected with human CD164 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram), and Isotype antibody on Expi293 transfected with irrelevant protein (Orange histogram).