

## 12-9455: Anti-APCDD1 antibody(DMC479); IgG1 Chimeric mAb

|                           |  |
|---------------------------|--|
| <b>Clonality :</b>        | Monoclonal                             |
| <b>Clone Name :</b>       | DMC479                                 |
| <b>Application :</b>      | FACS                                   |
| <b>Reactivity :</b>       | Human                                  |
| <b>Uniprot ID :</b>       | Q8J025                                 |
| <b>Alternative Name :</b> | B7323; DRAPC1; FP7019; HHS; HTS; HYPT1 |
| <b>Isotype :</b>          | Rabbit/Human Fc chimeric IgG1          |

### Description

Description :Anti-APCDD1 antibody(DMC479); IgG1 Chimeric mAb

This locus encodes an inhibitor of the Wnt signaling pathway. Mutations at this locus have been associated with hereditary hypotrichosis simplex. Increased expression of this gene may also be associated with colorectal carcinogenesis.[provided by RefSeq; Sep 2010]

### Product Info

|                            |   |
|----------------------------|---|
| <b>Amount :</b>            | 10 µg / 100 µg  |
| <b>Purification :</b>      | Purified from cell culture supernatant by affinity chromatography   |
| <b>Content :</b>           | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| <b>Storage condition :</b> | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |

### Application Note

FACS 1:100




Figure 1. Flow cytometry analysis with Anti-APCDD1 (DMC479) on Expi293 cells transfected with human APCDD1 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).