9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

12-9481: Anti-CD166 antibody(DMC682); IgG1 Chimeric mAb

Clonality :	Monoclonal
Clone Name :	DMC682
Application :	FACS
Reactivity :	Human
Uniprot ID :	Q13740-1
Alternative Name :	CD166; MEMD
Isotype :	Rabbit/Human Fc chimeric IgG1

Description

Description :Anti-CD166 antibody(DMC682); IgG1 Chimeric mAb
This gene encodes activated leukocyte cell adhesion molecule (ALCAM); also known as CD166 (cluster of differentiation 166); which is a member of a subfamily of immunoglobulin receptors with five immunoglobulin-like domains (VVC2C2C2) in the extracellular domain. This protein binds to T-cell differentiation antigene CD6; and is implicated in the processes of cell adhesion and migration. Multiple alternatively spliced transcript variants encoding different isoforms have been found. [provided by RefSeq; Aug 2011]

Product Info

Amount :	10 µg / 100 µg
Purification :	Purified from cell culture supernatant by affinity chromatography
Content :	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage condition :	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Application Note

FACS 1:100




Figure 1. CD166 protein is expressed on the surface of Expi293 cell membrane. Flow cytometry analysis with Anti-CD166 (DMC682) on Expi293 cells transfected with human CD166 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram), and Isotype antibody on Expi293 transfected with irrelevant protein (Orange histogram).

For Research Use Only. Not for use in diagnostic/therapeutics procedures.