

### 30-1374PE: PE Conjugated Anti-Human IgG (Fc) Monoclonal Antibody (Clone: EM-07)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	EM-07
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	PE
<b>Alternative Name :</b>	Immunoglobulin G Fc fragment
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Fusion protein of human IgG Fc fragment.

#### Description

Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the total plasma immunoglobulins. IgG has the highest stability of blood immunoglobulins ( $T_{1/2} = 21$  days) and is able of placental transfer. IgG is secreted by plasma cells at a comparably high rate as other immunoglobulins.

Specificity: The mouse monoclonal antibody EM-07 reacts with Fc part of human IgG heavy chain and with isolated Fc fragments.

#### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
<b>Content :</b>	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

#### Application Note

Flow cytometry: Recommended dilution: 1-5  $\mu\text{g/ml}$ .




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human IgG (Fc) (EM-07) PE antibody (concentration in sample 3  $\mu\text{g/ml}$ ).




Figure 2: Separation of human monocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human IgG (Fc) (EM-07) PE antibody (concentration in sample 3 µg/ml).