

12-9452B: Biotinylated Anti-CD164 antibody(DMC476); IgG1 Chimeric mAb

| | |
|---------------------------|------------------------------------|
| Clonality : | Monoclonal |
| Clone Name : | DMC476 |
| Application : | Flow Cyt |
| Reactivity : | Human |
| Gene : | CD164 |
| Uniprot ID : | Q04900 |
| Alternative Name : | LMOR; M-OR-1; MOP; MOR; MOR1; OPRM |
| Isotype : | IgG1 |

Description

This gene encodes one of at least three opioid receptors in humans; the mu opioid receptor (MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as beta-endorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse; such as nicotine; cocaine; and alcohol via its modulation of the dopamine system. The NM_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6 transmembrane domains. [provided by RefSeq; Oct 2013]

Product Info

| | |
|----------------------------|--|
| Amount : | 10 µg / 100 µg |
| Purification : | Purified from cell culture supernatant by affinity chromatography |
| Content : | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage condition : | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |

Application Note

Flow Cyt 1:100