

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-9613: Anti-CEACAM5 antibody(1B5), IgG1 Chimeric mAb

Clonality: Monoclonal

Clone Name: 1B5
Application: WB
Reactivity: Human
Gene: CEACAM5
Uniprot ID: P06731

Alternative Name: CEACAM-5;CD66e;CEA;Meconium antigen 100

Isotype: Rabbit/Human Fc chimeric IgG1

Description

This gene encodes a cell surface glycoprotein that represents the founding member of the carcinoembryonic antigen (CEA) family of proteins. The encoded protein is used as a clinical biomarker for gastrointestinal cancers and may promote tumor development through its role as a cell adhesion molecule. Additionally, the encoded protein may regulate differentiation, apoptosis, and cell polarity. This gene is present in a CEA family gene cluster on chromosome 19. Alternative splicing results in multiple transcript variants.

Product Info

Amount : $10 \mu g / 100 \mu g$

Purification : Purified from cell culture supernatant by affinity chromatography

Content: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.


Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

Storage condition: for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Application Note

WB 1:1000



"Figure 1.Anti-CEACAM5 antibody (Cat# 12-9613) at 1/1000 dilution Lane : HT55, whole cell lysate Secondary : Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution Predicted band size: 76kDa Observed band size: 50∏250 kDa"



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2. Flow cytometry analysis of antigen binding of anti-human CEACAM5 mAb(12-9613). (A)12-9613 does not bind to CHO-S cells that do not express CEACAM5. (B) A clear peak shift of 12-9613 was seen compared to the control when incubated with CEACAM5-expressing MKN45 cells, indicating strong binding of 12-9613 to CEACAM5. Antibodies were incubated at 5 $\mu g/mL$.