

### 36-1798: Monoclonal Antibody to VEGF (Vascular Endothelial Growth Factor)(Clone : SPM225)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPM225
<b>Application :</b>	IF,IHC
<b>Reactivity :</b>	Human, Mouse, Rat
<b>Gene :</b>	VEGFA
<b>Gene ID :</b>	7422
<b>Uniprot ID :</b>	P15692
<b>Format :</b>	Purified
<b>Alternative Name :</b>	VEGFA,VEGF
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Human VEGF189 recombinant protein

#### Description


This MAb recognizes proteins of 19-22kDa (reducing) and 38kDa-44kDa (non-reducing), identified as various isoforms of Vascular Endothelial Growth Factor or Vascular Permeability Factor (VEGF/VPF). It is highly specific to VEGF, which is a homodimeric, disulfide-linked glycoprotein with a close homology to platelet derived growth factor (PDGF). There are multiple isoforms of VEGF containing 206-, 189-, 165-, and 121-amino acid residues. The smaller two isoforms, VEGF165 and VEGF121, are secreted proteins and act as diffusible agents, whereas the larger two remain cell associated. VEGF/VPF plays an important role in angiogenesis, which promotes tumor progression and metastasis.

#### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

#### Application Note

Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (4-8ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA buffer, pH 8.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes)



Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with VEGF Monoclonal Antibody (SPM225).