

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-1798: Monoclonal Antibody to VEGF (Vascular Endothelial Growth Factor)(Clone: SPM225)

Clone Name : Monoclonal
Clone Name : SPM225
Application : IF,IHC

Reactivity: Human, Mouse, Rat

 Gene :
 VEGFA

 Gene ID :
 7422

 Uniprot ID :
 P15692

 Format :
 Purified

 Alternative Name :
 VEGFA,VEGF

 Isotype :
 Mouse IgG1, kappa

Immunogen Information: Human VEGF189 recombinant protein

## **Description**

This MAb recognizes proteins of 19-22kDa (reducing) and 38kDa-44kDa (non-reducing), identified as various isoforms of Vascular Endothelial Growth Factor or Vascular Permeability Factor (VEGF/VPF). It is highly specific to VEGF, which is a homodimeric, disulfide-linked glycoprotein with a close homology to platelet derived growth factor (PDGF). There are multiple isoforms of VEGF containing 206-, 189-, 165-, and 121-amino acid residues. The smaller two isoforms, VEGF165 and VEGF121, are secreted proteins and act as diffusible agents, whereas the larger two remain cell associated. VEGF/VPF plays an important role in angiogenesis, which promotes tumor progression and metastasis.

## **Product Info**

**Amount**: 100 μg

**Purification:** Affinity Chromatography

Content: 100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly

toxic.

Storage condition:


Storage condition:

Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (4-8ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA buffer, pH 8.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes)



Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with VEGF Monoclonal Antibody (SPM225).