

## 12-90418: Anti-ACVR2A antibody(15D6), IgG1 Chimeric mAb

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	15D6
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	ACVR2A
<b>Uniprot ID :</b>	P27037
<b>Alternative Name :</b>	ACVR2; ACTRII
<b>Isotype :</b>	Rabbit/Human Fc chimeric IgG1

### Description

This gene encodes a receptor that mediates the functions of activins, which are members of the transforming growth factor-beta (TGF-beta) superfamily involved in diverse biological processes. The encoded protein is a transmembrane serine-threonine kinase receptor which mediates signaling by forming heterodimeric complexes with various combinations of type I and type II receptors and ligands in a cell-specific manner. The encoded type II receptor is primarily involved in ligand-binding and includes an extracellular ligand-binding domain, a transmembrane domain and a cytoplasmic serine-threonine kinase domain. This gene may be associated with susceptibility to preeclampsia, a pregnancy-related disease which can result in maternal and fetal morbidity and mortality. Alternative splicing results in multiple transcript variants of this gene. [provided by RefSeq, Jun 2013]

### Product Info

<b>Amount :</b>	100 µg / 10 µg
<b>Purification :</b>	Purified from cell culture supernatant by affinity chromatography
<b>Content :</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % ? 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage condition :</b>	Store at -20C to -80C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

### Application Note

FACS 1/100




Figure 1. Flow cytometry analysis with 1 µg/mL Anti-ACVR2A(15D6) mAb on CHO-S cells(A) and Hu\_ACVR2A CHO-S Cell Line(B).