

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-18583: Cynomolgus CD142 Protein, His Tag

Gene: CD142 **Uniprot ID:** A0A2K5VX02

Alternative Name: TF; TFA; F3, Recombinant Cynomolgus CD142 protein with C-terminal 10? His tag

Description

This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces, for example, on monocytes. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. Platelets and monocytes have been shown to express this coagulation factor under procoagulatory and proinflammatory stimuli, and a major role in HIV-associated coagulopathy has been described. Platelet-dependent monocyte expression of coagulation factor III has been described to be associated with Coronavirus Disease 2019 (COVID-19) severity and mortality. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Aug 2020]

Molecular Weight: The protein has a predicted molecular mass of 26.4 kDa after removal of the signal peptide. The apparent molecular mass of cCD142-His is approximately 35-55 kDa due to glycosylation.

Tag:C-10?His tag

Product Info

Amount: 50μg / 10μg

Purification: The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

staining.

Content: Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

Storage condition : for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.




Figure 1. Cynomolgus CD142 Protein, His Tag on SDS-PAGE under reducing condition.