

12-8200: Anti-Chikungunya E1 Protein-Purified in vivo PLATINUM™ Functional Grade

Clonality :	Monoclonal
Clone Name :	CHK-166
Reactivity :	Mouse
Alternative Name :	CHIKV, Chikungunya virus, VLPs, Chikungunya virus-like particles

Description

Specificity: CHK-166 activity is directed against CHIKV E1.

Antigen Distribution: E1 is expressed on the surface of CHIKV.

Background: Chikungunya virus (CHIKV) is a mosquito-transmitted alphavirus that causes epidemics globally and has been declared a notable disease by the CDC^{1,2}. CHIKV is an enveloped virus with an 11.8-kb single-stranded, positive-sense RNA genome with two open reading frames^{3,4}. There are three main genotypes, having 95.2 to 99.8% amino acid identity: Asian, West African, and East/Central/South African (ECSA). The mature CHIKV virion is comprised of a nucleocapsid protein C and two glycoproteins, E1 and E2⁵. E1 participates in virus fusion. E2 functions in attachment to cells. E1 and E2 form 80 trimeric spikes on the virus surface⁶. CHK-166 is a neutralizing monoclonal antibody (MAb) that provides complete protection against lethality as prophylaxis in *Irf1^{+/+}* mice⁵. It was generated by infecting adult *Irf1^{+/+}* C57BL/6 mice with the La Reunion 2006 OPY-1 strain of CHIKV (CHIKV-LR) and boosting with recombinant CHIKV E2 protein or infectious CHIKV-LR. Myeloma cell-splenocyte fusions were screened for binding to CHIKV-LR infected cells and the resulting MAb was cloned for analysis. Neutralization escape variants were generated to map the CHK-166 epitope⁵. CHK-166 recognizes amino acids on domain II of E1, adjacent to the conserved fusion loop. All escape mutants had a single K61T mutation in the E1 protein. CHK-166 inhibits CHIKV infection in cell culture in a post-attachment neutralization assay⁵. CHK-166 also protects 63% of mice from death when a single dose is administered 24 h after CHIKV infection. If both CHK-166 and CHK-152 are administered post-infection in mice, then viral resistance is prevented and the treatment window is extended⁵. Additionally, combination CHK-152/CHK-166 MAb therapy in rhesus macaques reduces viral infection and spread, neutralizes reservoirs of infectious virus, and does not produce escape viruses⁷.

Product Info

Amount :	1 mg / 5 mg Purity: $\geq 98\%$ monomer by analytical SEC
Purification :	Preparation: Functional grade preclinical antibodies are manufactured in an animal free facility using in vitro cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates. Concentration: ≥ 5.0 mg/ml
Content :	Formulation: This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added. Due to inherent biochemical properties of antibodies, certain products may be prone to precipitation over time. Precipitation may be removed by aseptic centrifugation and/or filtration.
Storage condition :	This antibody may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at $\leq -70^\circ\text{C}$. Avoid Repeated Freeze Thaw Cycles.

Application Note

N ELISA FC