

32-1020: Acrp30 Tri Recombinant Protein

Alternative Name : Acrp30,AdipoQ,GBP-28,APM-1,ACDC.

Description

Source : HEK293 (Human embryonic kidney cell line). Trimeric form of Adiponectin Human trimeric form was expressed in HEK293 cells. The cysteine 39 was replaced with Alanine (C39A) 9. hAd-C39A can only form a trimer, but not a hexamer or an HMW form. Adiponectin is a hormone exclusively expressed from adipose tissue. Many studies demonstrate that Adiponectin has direct anti-diabetic, anti-atherogenic and anti-inflammatory functions. APM-1 can increase insulin sensitivity of skeletal muscle. Attenuate hepatic lipogenesis and gluconeogenesis, regulate NO production in endothelial cells, inhibit proliferation of smooth muscle cells and prevent lipid accumulation of macrophage cells. In the circulation, Adiponectin is present as three different oligomeric complexes, including the high molecular weight (HMW), the middle molecular weight (MMW, also called hexamer) and low molecular weight (MMW, also called trimer) forms 8. Different oligomeric complex of Adiponectin activates different signaling pathways and exerts distinct functions.

Product Info

Amount :	10 µg
Purification :	Greater than 95% as determined by SDS-PAGE.
Content :	Filtered (0.4µm) and lyophilized from 0.5 mg/ml in 0.05M phosphate buffer, 0.05M NaCl, pH 7.4.
Storage condition :	Store lyophilized Adiponectin at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.
Amino Acid :	ETTTQGPGLV LPLPKGAATG WMAGIPGHPG HNGAPGRDGR DGTPGEKGEK GDPGLIGPKG DIGETGVPGA EGPRGFPGIQ GRKGEPGEA YVYRSAFSVG LETYVTIPNM PIRFTKIFYN QQNHYDGSTG KFHCNIPGLY YFAYHIVYMK DVKVSLEFKD KAMLFYTDQY QENNVDQASG SVLLHLEVGQ QVWLQVYGEG ERNGLYADND NDSTFTGFL YHDTNDYKDD DDK.

Application Note

Add deionized water to a working volume of 0.5mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture. ED50= 3-8.5 Åµg/ml, as determined by its ability to inhibit proliferation of HASMCs induced by HB EGF.

