

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1509-B: Anti-CD195 / CCR5 Monoclonal Antibody (Clone:T21/8) Biotin Conjugated

Clonality: Monoclonal **Clone Name:** T21/8 ICC.FACS Application: Reactivity: Human Conjugate: Biotin Gene: CCR5 Gene ID: 1234 **Uniprot ID:** P51681 CCR5,CMKBR5 **Alternative Name:**

Immunogen Information: CCR5 peptide (Met1-Lys22) KLH conjugate

Mouse IgG1 kappa

Description

Isotype:

Specificity: The mouse monoclonal antibody T21/8 recognizes an extracellular epitope on the N-teminus of CD195, an approximately 45 kDa G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.

Description: CD195 / CCR5 (also known as CKR-5) is a receptor for inflammatory CC-chemokines (characterized by a pair of adjacent cysteine residues), such as MIP-1 alpha, MIP-1 beta, or RANTES. It is a G protein-associated seven-pass transmembrane protein expressed on resting T cells with memory/effector phenotype, monocytes, macrophages and immature dendritic cells. This chemokine receptor regulates the activation and directed migration of leukocytes. Importantly, along with CD4, CD195 / CCR5 functions as a major receptor for HIV. Their ligand is the viral glycoprotein gp120.

Product Info

Amount: 0.1 mg

Purification : Purified antibody is conjugated with biotin LC-NHS ester under optimum conditions and

unconjugated antibody and free biotin are removed by size-exclusion chromatography.

Content: Formulation: Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow cytometry: Recommended dilution: 1-5 µg/ml



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD195 (T21/8) biotin antibody (concentration in sample 5 μ g/ml, Streptavidin APC).




Figure 2: Separation of human monocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD195 (T21/8) biotin antibody (concentration in sample 5 μ g/ml, Streptavidin APC).