

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

19-10001: Purified anti-CD34 Antibody (Clone: QBEnd/10)

Clonality: Monoclonal
Clone Name: QBEnd/10
Application: IHC
Reactivity: Human
Gene: CD34
Gene ID: 947
Format: Purified

Alternative Name: Hematopoietic progenitor cell antigen CD34, CD34 antigen

Isotype : Mouse IgG1, κ

Immunogen Information: This antibody was generated using a vesicular suspension of a perfusate of human term

placenta

Description

CD34, also known as gp105-120, is a type I monomeric sialomucin-like glycophosphoprotein with an approximate molecular weight of 105-120 kD. Selectively expressed on the majority of hematopoietic stem/progenitor cells, bone marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nervous tissue, CD34 is a commonly used marker to identify human hematopoietic stem/progenitor cells. According to the differential sensitivity to enzymatic cleavage, four groups of epitopes of CD34 have been described. CD34 mediates cell adhesion and lymphocytes homing through binding to L-selectin and E-selectin ligands.

Product Info

Amount: 50ul

Purification : The antibody was purified by affinity chromatography.

Content: Concentration: 1.0 mg/ml

Formulation: Phosphate-buffered solution with BSA + 0.09% NaN3.

Storage condition : Store between 2°C and 8°C.

Application Note

This antibody is effective in immunohistochemistry (IHC).

This antibody is effective in immunohistochemistry (IHC).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Robust cytoplasmic and membrane staining of endothelial cells in frozen human placenta with antibody CD34.




Figure 2: SeqIF[™] (sequential immunofluorescence) staining on COMET[™] of Purified anti-CD34 (clone QBEnd/10, yellow) on formalin-fixed paraffin-embedded human pancreatic carcinoma tissue at 1.25 μ g/mL. Alexa Fluor[™] Plus 647 Goat anti-Mouse IgG antibody was used as a secondary antibody. Nuclei were counterstained with DAPI (blue). Tissue underwent an all-in-one dewaxing and antigen retrieval preprocessing.