

30-2818-AC: Anti-Hu CD199 APC

Clonality : Monoclonal
Clone Name : C9Mab-1
Application : FACS
Reactivity : Human
Conjugate : APC
Gene : CCR9
Gene ID : 10803
Uniprot ID : P51686
Alternative Name : C-C motif chemokine receptor 9 GPR-9-6, GPR28, CCR9
Isotype : Mouse IgG1 kappa
Immunogen Information : CD199 transfected CHO cells

Description

CD199 (CCR9) is a G-protein-coupled 7 TM chemokine receptor for TECK (SCYA25) chemokine. It is expressed strongly in thymus, at lower level in bone marrow and spleen, as well as on a subset of memory T cells specialized for mucosal homing. CD199 appears to confer homing properties to the small intestine on memory T cells. On the other hand it functions as a coreceptor for HIV-1.

Specificity : The mouse monoclonal antibody C9Mab-1 recognizes an extracellular epitope of CD199, a 7-transmembrane chemokine receptor.

Product Info

Amount : 100 tests (T100)
Purification : Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Content : Storage Buffer: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage condition : Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1: Separation of MOLT-4 cells stained using anti-human CD199 (C9Mab-1) APC antibody (concentration in sample 1.67 µg/ml, red-filled) from MOLT-4 cells stained using mouse IgG1 isotype control (MOPC-21) APC antibody (concentration in sample 1.67 µg/ml, same as CD199 APC concentration, black-dashed) in flow cytometry analysis (surface staining).