

### 30-2711-AC: Anti-Hu TCR Vgamma9 APC

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	B3
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human,Non-Human Primates
<b>Conjugate :</b>	APC
<b>Gene ID :</b>	35
<b>Alternative Name :</b>	TCRVG9
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	human T cells

#### Description

TCR Vgamma9 is a variant of TCR gamma chain, that is present on a subset of human gamma/delta T cells. TCR Vgamma9/Vdelta2 T cells are able to recognize and kill various tumor cells, as this receptor heterodimer binds to certain phosphoantigens, expressed by tumors.

Specificity :The mouse monoclonal antibody B3 recognizes an extracellular epitope of human TCR Vgamma9, a variant of TCR gamma chain, which is expressed on a gamma/delta T cell subpopulation with cytolytic activity against various tumor cells.

#### Product Info

<b>Amount :</b>	100 tests (T100)
<b>Purification :</b>	Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
<b>Content :</b>	Storage Buffer: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

#### Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human TCR Vgamma9 (B3) APC antibody (concentration in sample 1.67 µg/ml).




Figure 2: Flow cytometry multicolor surface staining pattern of human lymphocytes stained using anti-human CD3 (UCHT1) PE antibody (20  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood) and anti-human TCR Vgamma9 (B3) APC antibody (concentration in sample 1.67  $\mu$ g/ml).




Figure 3: Separation of human CD3 positive TCR Vgamma9 positive T cells (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human TCR Vgamma9 (B3) APC antibody (concentration in sample 1.67  $\mu$ g/ml).