

30-2711-FITC: Anti-Hu TCR Vgamma9 FITC

Clonality : Monoclonal
Clone Name : B3
Application : FACS
Reactivity : Human, Non-Human Primates
Conjugate : FITC
Gene ID : 35
Alternative Name : TCRVG9
Isotype : Mouse IgG1
Immunogen Information : human T cells

Description

TCR Vgamma9 is a variant of TCR gamma chain, that is present on a subset of human gamma/delta T cells. TCR Vgamma9/Vdelta2 T cells are able to recognize and kill various tumor cells, as this receptor heterodimer binds to certain phosphoantigens, expressed by tumors.

Specificity : The mouse monoclonal antibody B3 recognizes an extracellular epitope of human TCR Vgamma9, a variant of TCR gamma chain, which is expressed on a gamma/delta T cell subpopulation with cytolytic activity against various tumor cells.

Product Info

Amount : 100 tests (T100)
Purification : Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Content : Storage Buffer: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage condition : Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 4 μ l reagent / 100 μ l of whole blood or 10^6 cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.




Figure 1: Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human TCR Vgamma9 (B3) FITC antibody (concentration in sample 0.56 μ g/ml).




Figure 2: Flow cytometry multicolor surface staining pattern of human lymphocytes stained using anti-human CD3 (UCHT1) APC antibody (10 μ l reagent / 100 μ l of peripheral whole blood) and anti-human TCR Vgamma9 (B3) FITC antibody (concentration in sample 0.56 μ g/ml).




Figure 3: Separation of human CD3 positive TCR Vgamma9 positive T cells (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human TCR Vgamma9 (B3) APC antibody (concentration in sample 0.56 μ g/ml).