w abeomics

32-1041: APOA5 HEK Recombinant Protein

Alternative Name : Apolipoprotein A-V,Apo-AV,ApoA-V,Apolipoprotein A5,Regeneration-associated protein 3,APOA5,RAP3,APOAV.

Description

Source : HEK293 APOA5 Human Recombinant Flag-Tagged Fusion Protein is 40.1 kDa protein containing 354 amino acid residues of the APOA5 Human and 11 additional amino acid residues of flagTag. ApoA5 is an important regulator of serum triglyceride concentrations. An ApoA5 mouse nock-out model produced an approximately four fold increase in serum triglycerides, whereas a knock-in model human ApoA5 produced 50-70% lower concentrations of mouse serum triglycerides. Furthermore, peroxisome proliferators-activated receptoragonists, which are used clinically to lower serum triglyceride concentrations, cause increased ApoA5 mRNA expression. There is very little known about ApoA5 protein in human serum. Lately, it was established that ApoA5 is present in the human serum and it can be detected by polyclonal antibodies against both the HN2 and COOH termini, but at much lower concentration than other apolipoproteins.

Product Info

| Amount : | 10 µg |
|---------------------|--|
| Purification : | Greater than 70% as determined by SDS-PAGE. |
| Content : | APOA5 Human was filtered (0.4 μm) and lyophilized from 0.5 mg/ml supplied in 20mM TRIS and 50mM NaCl, pH 7.5 |
| Storage condition : | Store lyophilized APOA5 Human at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted APOA5 Human can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C. |
| Amino Acid : | RKGFWDYFSQ TSGDKGRVEQ IHQQKMAREP ATLKDSLEQD LNNMNKFLEK LRPLSGSEAP RLPQDPVGMR RQLQEELEEV KARLQPYMAE AHELVGWNLE GLRQQLKPYT MDLMEQVALR VQELQEQLRV VGEDTKAQLL GGVDEAWALL QGLQSRVVHH TGRFKELFHP YAESLVSGIG RHVQELHRSV APHAPASPAR LSRCVQVLSR KLTLKAKALH ARIQQNLDQL REELSRAFAG TGTEEGAGPD PQMLSEEVRQ RLQAFRQDTY LQIAAFTRAI DQETEEVQQQ LAPPPPGHSA FAPEFQQTDS GKVLSKLQAR LDDLWEDITH SLHDQGHSHL GDPAAADYKD DDDK |

Application Note

It is recommended to add deionized water to prepare a working stock solution of approximately 0.5mg/ml and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it on cell culture.



For Research Use Only. Not for use in diagnostic/therapeutics procedures.