9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

12-9213-PE: PE-conjugated Anti-CD160 antibody(DMC270); IgG1 Chimeric mAb

Clonality :	Monoclonal
Clone Name :	DMC270
Application :	Flow Cyt
Reactivity :	Human
Conjugate :	PE-conjugated
Gene :	CD160
Uniprot ID :	O95971
Alternative Name :	BY55; NK1; NK28
Isotype :	Rabbit/Human Fc chimeric IgG1

Description

CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich; glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single Ig-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells; spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16 cells whereas among circulating T cells its expression is mainly restricted to TCRgd bearing cells and to TCRab CD8brightCD95 CD56 CD28-CD27- cells. In tissues; CD160 is expressed on all intestinal intraepithelial lymphocytes. CD160 shows a broad specificity for binding to both classical and nonclassical MHC class I molecules.

Product Info

Amount :	100 Test
Purification :	Purified from cell culture supernatant by affinity chromatography
Content :	Liquid PBS with 0.05% Proclin300, 1% BSA
Storage condition :	Store at 2°C-8°C for 6 months

Application Note

Flow Cyt 1:100

For Research Use Only. Not for use in diagnostic/therapeutics procedures.