

**12-9428-PE: PE-conjugated Anti-UPA antibody(DMC440); IgG1 Chimeric mAb**

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	DMC440
<b>Application :</b>	Flow Cyt
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	PE-conjugated
<b>Gene :</b>	UPA
<b>Uniprot ID :</b>	P00749
<b>Alternative Name :</b>	ATF; BDPLT5; QPD; u-PA; UPA; URK
<b>Isotype :</b>	Rabbit/Human Fc chimeric IgG1

**Description**

This gene encodes a secreted serine protease that converts plasminogen to plasmin. The encoded preproprotein is proteolytically processed to generate A and B polypeptide chains. These chains associate via a single disulfide bond to form the catalytically inactive high molecular weight urokinase-type plasminogen activator (HMW-uPA). HMW-uPA can be further processed into the catalytically active low molecular weight urokinase-type plasminogen activator (LMW-uPA). This low molecular weight form does not bind to the urokinase-type plasminogen activator receptor. Mutations in this gene may be associated with Quebec platelet disorder and late-onset Alzheimer's disease. Alternative splicing results in multiple transcript variants; at least one of which encodes an isoform that is proteolytically processed.

**Product Info**

<b>Amount :</b>	100 Test
<b>Purification :</b>	Purified from cell culture supernatant by affinity chromatography
<b>Content :</b>	Liquid PBS with 0.05% Proclin300, 1% BSA
<b>Storage condition :</b>	Store at 2°C-8°C for 6 months

**Application Note**

Flow Cyt 1:100