

## 12-9434-PE: PE-conjugated Anti-CD117 antibody(DMC446), IgG1 Chimeric mAb

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	DMC446
<b>Application :</b>	Flow Cyt
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	PE-conjugated
<b>Gene :</b>	CD117
<b>Uniprot ID :</b>	P10721
<b>Alternative Name :</b>	C-Kit; CD117; MASTC; PBT; SCFR; KIT
<b>Isotype :</b>	Rabbit/Human Fc chimeric IgG1

### Description

This gene encodes a receptor tyrosine kinase. This gene was initially identified as a homolog of the feline sarcoma viral oncogene v-kit and is often referred to as proto-oncogene c-Kit. The canonical form of this glycosylated transmembrane protein has an N-terminal extracellular region with five immunoglobulin-like domains; a transmembrane region; and an intracellular tyrosine kinase domain at the C-terminus. Upon activation by its cytokine ligand; stem cell factor (SCF); this protein phosphorylates multiple intracellular proteins that play a role in in the proliferation; differentiation; migration and apoptosis of many cell types and thereby plays an important role in hematopoiesis; stem cell maintenance; gametogenesis; melanogenesis; and in mast cell development; migration and function. This protein can be a membrane-bound or soluble protein. Mutations in this gene are associated with gastrointestinal stromal tumors; mast cell disease; acute myelogenous leukemia; and piebaldism. Multiple transcript variants encoding different isoforms have been found for this gene.

### Product Info

<b>Amount :</b>	100 Test
<b>Purification :</b>	Purified from cell culture supernatant by affinity chromatography
<b>Content :</b>	Liquid PBS with 0.05% Proclin300, 1% BSA
<b>Storage condition :</b>	Store at 2°C-8°C for 6 months

### Application Note

Flow Cyt 1:100