

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1079: BMP 2 Recombinant Protein

Alternative Name: BMP-2,BMP2A.

Description

Source: Escherichia Coli. Bone Morphogenetic Protein-2 Human Recombinant produced in E.Coli is a homodimeric, non-glycosylated polypeptide chain containing 2x115 amino acids and having a molecular mass of 26kDa. The BMP-2 is purified by proprietary chromatographic techniques. BMP2 belongs to the transforming growth factor-beta (TGFB) superfamily. Bone morphogenic protein induces bone formation. BMP2 is a candidate gene for the autosomal dominant disease of fibrodysplasia (myositis) ossificans progressiva.

Product Info

Amount: 10 μg

Purification: Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

BMP2 was lyophilized from a concentrated (1mg/ml) sterile solution containing 10mM sodium

Content:

BMP2 was lyophilized from a concentrated (1mg/ml) sterile solution containing 10mM sodium

citrate pH=3.5.

Lyophilized Bone Morphogenetic Protein-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution BMP2 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add

a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: MQAKHKQRKR LKSSCKRHPL YVDFSDVGWN DWIVAPPGYH AFYCHGECPF PLADHLNSTN

HAIVQTLVNS VNSKIPKACC VPTELSAISM LYLDENEKVV LKNYQDMVVE GCGCR.

Application Note

It is recommended to reconstitute the lyophilized Bone Morphogenetic Protein-2 in sterile 20mM AcOH (acetic Acid) not less than $100\text{Å}\mu\text{g/ml}$, which can then be further diluted to other aqueous solutions. The ED50 as determined by its ability to induce alkaline phosphatase production by ATDC-5 cells is 0.5-1.0 $\text{Å}\mu\text{g/ml}$.

