

## 12-90482: Anti-CXCL5 antibody(3C1), Rabbit mAb

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	3C1
<b>Application :</b>	ELISA
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	Unconjugated
<b>Gene :</b>	CXCL5
<b>Uniprot ID :</b>	P42830
<b>Alternative Name :</b>	ENA-78;SCYB5
<b>Isotype :</b>	Rabbit mAb

### Description

This gene encodes a protein that is a member of the CXC subfamily of chemokines. Chemokines, which recruit and activate leukocytes, are classified by function (inflammatory or homeostatic) or by structure. This protein is proposed to bind the G-protein coupled receptor chemokine (C-X-C motif) receptor 2 to recruit neutrophils, to promote angiogenesis and to remodel connective tissues. This protein is thought to play a role in cancer cell proliferation, migration, and invasion. [provided by RefSeq, May 2013]

### Product Info

<b>Amount :</b>	100 µg / 10 µg
<b>Purification :</b>	Purified from cell culture supernatant by affinity chromatography
<b>Content :</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization.
<b>Storage condition :</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

### Application Note

ELISA 1/5000-10000




Figure 1. ELISA plate pre-coated by 1 µg/ml (100 µl/well) Human CXCL5 Protein can bind Rabbit anti-CXCL5 monoclonal antibody(clone: 3C1) in a linear range of 1-10 ng/ml.