

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-1151: EPO a Recombinant Protein

**Alternative Name:** Erythropoietin-Alpha, EPO-a, EPO-alpha, Epoetin, EP, MGC138142.

## **Description**

Source: Chinese Hamster Ovary Cells(CHO). Erythropoietin-alpha Human Recombinant is produced in Chinese hamster ovary (CHO) cells by recombinant DNA technology is a single, polypeptide chain containing 166 amino acids and having a predicted molecular mass of 21,000 Dalton and apparent glycosylated molecular mass of 36-40kDa. EPO-a is purified by proprietary chromatographic techniques. This gene is a member of the EPO/TPO family and encodes a secreted, glycosylated cytokine composed of four alpha helical bundles. The protein is found in the plasma and regulates red cell production by promoting erythroid differentiation and initiating hemoglobin synthesis. This protein also has neuroprotective activity against a variety of potential brain injuries and antiapoptotic functions in several tissue types.

## **Product Info**

Amount: 50 µg

**Purification:** Greater than 98.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Each mg of lyophilized powder contains 0.59 mg sodium citrate, 0.58 mg sodium chloride and Content:

0.006 mg citric acid.

Lyophilized Erythropoietin-a although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution EPO-alpha should be stored at 4°C between 2-7 Storage condition:

days and for future use below -18°C. For long term storage it is recommended to add a carrier

protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

APPRLICDSR VLERYLLEAK EAENITTGCA EHCSLNENIT VPDTKVNFYA WKRMEVGQQA **Amino Acid:** 

VEVWQGLALL SEAVLRGQAL LVNSSQPWEP LQLHVDKAVS GLRSLTTLLR ALGAQKEAIS

PPDAASAAPL RTITADTFRK LFRVYSNFLR GKLKLYTGEA CRTGDR.

## **Application Note**

It is recommended to reconstitute the lyophilized Epoetin-a in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The Specific Activity was measured by Normocyth -aemic mice and was found to be 150,000 IU/mg.

