

36-3827: Recombinant Monoclonal Antibody to TRIM29 (Lung Squamous Cell Carcinoma Marker)(Clone : TRIM29/9256R)

Clonality :	Monoclonal
Clone Name :	TRIM29/9256R
Application :	IHC,WB
Reactivity :	Human
Gene :	TRIM29
Gene ID :	23650
Uniprot ID :	Q14134
Format :	Purified
Alternative Name :	Tripartite motif-containing protein 29, Ataxia telangiectasia group D-associated protein, Ataxia telangiectasia group D complementing gene (ATDC); Tripartite motif-containing protein 29 (TRIM29);
Isotype :	IgG / Kappa
Immunogen Information :	Recombinant fragment (around aa 1-200) of human TRIM29 protein (exact sequence is proprietary)

Description

It recognizes a 66kDa protein, which is identified as Tripartite motif-containing protein 29 (TRIM29). It interacts with the intermediate filament protein vimentin, a substrate for the PKC family of protein kinases, and with hPKCI-1, an inhibitor of the PKCs. TRIM29 protein contains both zinc finger and leucine zipper motifs, suggesting that the it may form homodimers and possibly associate with DNA. High expression of TRIM29 has been reported in gastric cancer and pancreatic cancer, and correlates with enhanced tumor growth and lymph node metastasis. TRIM29 is also able to distinguish lung squamous cell carcinoma from lung adenocarcinoma with ~90% positive accuracy, when used in a panel with TTF-1, p63, CK5/6, and Napsin-A antibodies.

Product Info

Amount :	20 µg / 100 µg
Purification :	Prepared in 1mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Content :	This antibody is available for research use only and is not approved for use in diagnosis. Supplied as 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Application Note

Immunohistochemistry (IHC)

1-2ug/ml

IHC Protocol

30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Western Blot (WB)

2-4ug/ml

Western Blot Protocol




Figure 1: Formalin-fixed, paraffin-embedded human esophagus stained with TRIM29 Recombinant Rabbit Monoclonal Antibody (TRIM29/9256R). Inset: PBS instead of primary antibody; secondary only negative control.




Figure 2: SDS-PAGE Analysis of Purified TRIM29 Recombinant Rabbit Monoclonal Antibody (TRIM29/9256R). Confirmation of Purity and Integrity of Antibody.




Figure 3: TRIM29 Antibody in Western Blot (WB) Western Blot analysis of A431 cell lysate using TRIM29 Recombinant Rabbit Monoclonal Antibody (TRIM29/9256R).