

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-1159: FGF 1 Sf9 Recombinant Protein

Alternative Name: HBGF-1,ECGF-beta,FIBP,FGFIBP,FIBP-1,ECGF,ECGFA,GLIO703,FGF1,FGF-a.

Description

Source: Baculovirus. Fibroblast Growth Factor-1 Human Recombinant (FGF-1) produced in Sf9 insect cells is a single, glycosylated, polypeptide chain containing 140 amino acids and having a molecular mass of 15803 Dalton. The FGF-a is purified by proprietary chromatographic techniques. Acidic fibroblast growth factor is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein functions as a modifier of endothelial cell migration and proliferation, as well as an angiogenic factor. It acts as a mitogen for a variety of mesoderm- and neuroectoderm-derived cells in vitro, thus is thought to be involved in organogenesis. Three alternatively spliced variants encoding different isoforms have been described. The heparin-binding growth factors are angiogenic agents in vivo and are potent mitogens for a variety of cell types in vitro. There are differences in the tissue distribution and concentration of these 2 growth factors.

Product Info

Amount: 10 µc

Purification: Greater than 95.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Fibroblast Growth Factor-acidic although stable at 4°C for 3 weeks, should be stored desiccated

Storage condition: below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Please prevent freeze-thaw cycles.

Amino Acid: The sequence of the first five N-terminal amino acids as determined and was found to be Met-

Phe-Asn-Leu-Pro.

Application Note

The ED50, calculated by the dose-dependant proliferation of BAF3 cells expressing FGF receptors (measured by 3H-thymidine uptake) is <10 ng/ml, corresponding to a specific activity of 100,000IU/mg.

