

32-1204: Flt3 Ligand Recombinant Protein

Alternative Name : Fms-related tyrosine kinase 3 ligand,FLK2,STK1,CD135,Stem Cell Tyrosine Kinase 1,FLT3LG,Flt3.

Description

Source : Escherichia Coli. Flt3-Ligand Human Recombinant produced in E.Coli is a non-glycosylated, polypeptide chain containing 155 amino acids and having a molecular mass of approximately 17.6kDa. Flt3-Ligand is purified by proprietary chromatographic techniques. FLT3 ligand is a receptor for the fl cytokine has a tyrosine-protein kinase activity & a growth factor that regulates proliferation of early hematopoietic cells. Flt3-Ligand synergizes with other CSFs and interleukins to induce growth and differentiation.

Product Info

Amount :	10 µg
Purification :	Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	The protein was lyophilized with no additives.
Storage condition :	Lyophilized Flt3-Ligand although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Flt3-Ligand should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid :	TQDCSFQHSP ISSDFAVKIR ELSDYLLQDY PVTVASNLQD EELCGGLWRL VLAQRWMERL KTVAGSKMQG LLERVNTEIH FVTKCAFQPP PSCLRFVQTN ISRLLQETSE QLVALKPWIT RQNFSRCLEL QCQPDSSTLP PPWSPRPLEA TAPTA.

Application Note

It is recommended to reconstitute the lyophilized Flt3-L in sterile 18M $\bar{\bar{A}}$ -cm H₂O not less than 100 $\bar{\bar{A}}$ µg/ml, which can then be further diluted to other aqueous solutions. The ED₅₀ as calculated by the dose-dependent stimulation of the proliferation of human AML5 cells is less than 1.0 ng/ml, corresponding to a Specific Activity of 1.0 $\bar{\bar{A}}$ 106 IU/mg.