

32-1254: G-CSF HEK Recombinant Protein

Alternative Name : CSF-3,MGI-1G,GM-CSF beta,Pluripoietin,Filgrastim,Lenograstim,G-CSF,MGC45931,GCSF.

Description

Source : HEK. G-CSF Human Recombinant produced in HEK cells is a glycosylated monomer, having a molecular weight range of 21-25kDa due to glycosylation. The G-CSF is purified by proprietary chromatographic techniques. GCSF is a cytokine that controls the production, differentiation, and function of granulocytes. The active protein is found extracellularly. Three transcript variants encoding three different isoforms have been found for the GCSF gene. Granulocyte/macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. This csf induces granulocytes.

Product Info

Amount :	10 µg
Purification :	Greater than 95% as observed by SDS-PAGE.
Content :	The G-CSF was lyophilized from 1mg/ml in 1xPBS.
Storage condition :	Lyophilized G-CSF although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution G-CSF should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Application Note

It is recommended to reconstitute the lyophilized G-CSF in sterile 1xPBS containing 0.1% endotoxin-free recombinant HSA. The specific activity was determined by the dose-dependent stimulation of the proliferation of murine M-NFS-60 cells (Mouse Myeloid Leukemia indicator cell line), the ED50 is <0.1ng/ml.

