

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-1265: chGH Recombinant Protein

**Alternative Name:** GH1,GH,GHN,GH-N,hGH-N,Pituitary growth hormone,Growth hormone 1,Somatotropin.

## **Description**

Source: Escherichia Coli. Growth Hormone Chicken Recombinant (cGH) produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 191 amino acids with an additional Ala at its N-terminus and having a molecular mass of 22255 Dalton. GH Chicken recombinant is purified by proprietary chromatographic techniques. GH is a member of the somatotropin/prolactin family of hormones which play an important role in growth control. The gene, along with four other related genes, is located at the growth hormone locus on chromosome 17 where they are interspersed in the same transcriptional orientation; an arrangement which is thought to have evolved by a series of gene duplications. The five genes share a remarkably high degree of sequence identity. Alternative splicing generates additional isoforms of each of the five growth hormones, leading to further diversity and potential for specialization. This particular family member is expressed in the pituitary but not in placental tissue as is the case for the other four genes in the growth hormone locus. Mutations in or deletions of the gene lead to growth hormone deficiency and short stature.

## **Product Info**

Amount: 100 µg

**Purification:** Greater than 99.0% as determined by:(a) Analysis by SDS-PAGE gel.(b) Analysis by SEC-HPLC.

The protein was lyophilized from a concentrated (1mg/ml) solution with 0.3% NaHCO3 adjusted Content:

to pH 8.

Lyophilized GH Chicken although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution at > 0.1 mg/ml and filter sterilization GH can be Storage condition:

stored at 4°C for several weeks. At lower concentration addition of a carrier protein (0.1% HSA

or BSA) is recommended. Please prevent freeze-thaw cycles.

Amino Acid: The sequence of the first five N-terminal amino acids was determined and was found to be Ala-

Thr-Phe-Pro-Ala.

## **Application Note**

It is recommended to reconstitute the lyophilized GH Chicken Recombinant in sterile water or 0.4% NaHCO3 adjusted to pH 8-9, not less than 100µg/ml and not more than 3 mg/ml, which can then be further diluted to other aqueous solutions, preferably in presence of carrier protein. GH Chicken Recombinantis fully biologically active in homologous assays and in PDF-P1 3B9 cells stably transfected with rabbit GH receptors.

